

Mitochondrial DNA m.3243A>G heteroplasmy affects multiple aging phenotypes and risk of mortality

Gregory J. Tranah ^{1*}, Shana M. Katzman ², Kevin Lauterjung ¹, Kristine Yaffe ³, Todd M. Manini ⁴, Stephen Kritchevsky ⁵, Anne B. Newman ⁶, Tamara B. Harris ⁷, and Steven R. Cummings ¹.

¹ California Pacific Medical Center Research Institute, San Francisco, CA, 94107, USA

² L.A. Eye Center and Clinic, Los Angeles, CA, 90037, USA

³ Departments of Psychiatry, Neurology, and Epidemiology, University of California, San Francisco and the San Francisco VA Medical Center, San Francisco, CA 94121, USA


⁴ Department of Aging and Geriatric Research, University of Florida, Gainesville, FL 32601, USA

⁵ Sticht Center on Aging, Wake Forest School of Medicine, Winston-Salem, NC, 27157, USA

⁶ Department of Epidemiology, University of Pittsburgh, Pittsburgh, PA, 15213, USA

⁷ Intramural Research Program, Laboratory of Epidemiology and Population Sciences, National Institute on Aging, Bethesda, MD, 20892, USA

Supplementary Figure 1. Mitochondrial m.3243A>G not associated association with cancer-related mortality. Survival was compared across tertiles of m.3243A>G heteroplasmy with a total of 263 participants included in each tertile. No significantly increased risks with cancer-related mortality were observed Analyses adjusted for age, sex, race, and clinic site.



Supplementary Figure 2. Mitochondrial m.3243A>G association with CVD-related mortality.

Survival was compared across tertiles of m.3243A>G heteroplasmy with a total of 263 participants included in each tertile. No significantly increased risks with CVD-related mortality were observed. Analyses adjusted for age, sex, race, and clinic site.

